

Diagram illustrating the structure of the human $\alpha 1(I)$ procollagen gene. The gene consists of three exons (EXON 1, EXON 2, EXON 3) and two introns (INTRON 1, INTRON 2). The sizes of the regions are indicated: EXON 1 (64 bp), EXON 2 (97 bp), EXON 3 (4.2 kb), INTRON 1 (2.4 kb), and INTRON 2 (14 kb). The 5' and 3' ends of the gene are also marked.

FIG. 1




FIG. 2A




FIG. 2B





FIG. 3

00816546-032604
FO922E0-9459T860



BamHI

BglII

EcoRI

HindIII

Xba I

FIG. 4

094646.02501
T09220" 94957860




Figure 5




Figure 6

Figure 7. Cell survival in electroporation of BFF transfected with pPNT and pPRP.






Figure 8. Electroporation of BFF transfected with pPNT - second experiment



0981545-022601
T09220-945T860

Figure 9. Electroporation of BFF transfected with pPRP.



T09220" 9459T860

Figure 10 G418 Treatment of untransfected BFF cells.




Figure 11





Figure 12. Genomic DNA organization of bovine PrP and the gene targeting strategy.



Top: The bovine PrP gene is composed of 3 exons and two introns, spanning an over 20 kb region [43]. Exon 1 and 2, which are 53 by and 98 by in size respectively, are transcribed as S' UTR, and the third exon contains sequences of 10 by S' UTR, 795 by coding region and about 3.3 kb 3' UTR. Intron 1 and 2 are about 2.4 kb and 14 kb in size.

Middle: The gene targeting vector contains a part of intron 2 sequence (at least 7 kb, exon 3 in which the bovine PrP coding sequence is completely deleted and replaced with promoterless neomycin resistant gene, and partial downstream genomic sequence of exon 3. The expression of neomycin resistant gene is under the control of the endogenous PrP promoter and its regulatory elements.

Bottom: The targeted bovine PrP allele after homologous recombination. Shaded boxes, exons; open boxes with names of gene, coding sequence the genes. ATP, start codon.